

Anomalous isotope effect near a 2.5 Lifshitz transition in a multi-band multi-condensate superconductor made of a superlattice of stripes

Andrea Perali ^{1,2}, Davide Innocenti ³, Antonio Valletta ⁴, Antonio Bianconi ^{2,5,6}

¹*School of Pharmacy, Physics Unit, University of Camerino, 62032 Camerino, Italy*

²*Mediterranean Institute of Fundamental Physics, Via Appia Nuova 31, 00040 Marino, Italy*

³*CNR-SPIN and Dipartimento di Ingegneria Informatica Sistemi e Produzione,*

"Tor Vergata" University of Rome, Via del Politecnico 1, 00133 Roma, Italy

⁴*CNR-IMM, Sezione di Roma, Via del Fosso del Cavaliere 100, 00133 Roma, Italy*

⁵*Rome International Center for Materials Science Superstripes (RICMASS) Via dei Sabelli 119A, 00185 Roma, Italy*

⁶*Department of Physics, Sapienza University of Rome, P. le A. Moro 2, 00185 Roma, Italy*

The doping dependent isotope effect on the critical temperature (T_c) is calculated for multi-band multi-condensate superconductivity near a 2.5 Lifshitz transition. We focus on multi-band effects that arises in nano-structures and in density wave metals (like spin density wave or charge density wave) as a result of the band folding. We consider a superlattice of quantum stripes with finite hopping between stripes near a 2.5 Lifshitz transition for appearing of a new sub-band making a circular electron-like Fermi surface pocket. We describe a particular type of BEC (Bose-Einstein Condensate) to BCS (Bardeen-Cooper-Schrieffer condensate) crossover in multi-band / multi-condensate superconductivity at a metal-to-metal transition that is quite different from the standard BEC-BCS crossover at an insulator-to-metal transition. The electron wave-functions are obtained by solving the Schrödinger equation for a one-dimensional modulated potential barrier. The k-dependent and energy dependent superconducting gaps are calculated using the k-dependent anisotropic Bardeen-Cooper-Schrieffer (BCS) multi-gap equations solved joint with the density equation, according with the Leggett approach currently used now in ultracold fermionic gases. The results show that the isotope coefficient strongly deviates from the standard BCS value 0.5, when the chemical potential is tuned at the 2.5 Lifshitz transition for the metal-to-metal transition. The critical temperature T_c shows a minimum due to the Fano antiresonance in the superconducting gaps and the isotope coefficient diverges at the point where a BEC coexists with a BCS condensate. On the contrary (T_c) reaches its maximum and the isotope coefficient vanishes at the crossover from a polaronic condensate to a BCS condensate in the new appearing sub-band.

PACS numbers: 74.62.-c, 74.70.Ad, 74.78.Fk

INTRODUCTION

Experiment. While for many years most of the mechanisms proposed for high temperature superconductivity have assumed a homogeneous lattice, recently new experimental results have shifted the theoretical research toward complex materials showing multi-band / multi-condensate superconductivity. Quantum oscillations have recently revealed the presence of one small electron pocket in the Fermi surface of cuprate superconductors related with the stripe phase [1–5]. It has also been proposed that the Fermi surface reconstruction in cuprates could arise from intrinsic effects in a doped Mott insulator [6–13]. These new results are leading the community to consider the new possibility of multi-band / multi-condensate superconductivity in charge density wave metals. The fundamental theoretical problems in this new scenario are similar to the superconductivity in ultra-narrow materials where the multi-band superconductivity is generated by quantum size effects due to the material lattice structure. Moreover the short range lattice structure in cuprates deviates from the simple average structure in the region of the phase diagram of cuprates $T_c < T < T^*$, where deviations from a Fermi liquid metallic state are sizeable. Experimental fast and

local structural methods, like Extended X-ray Absorption Fine Structure (EXAFS) [14], and X-ray Absorption Near Edge Structure (XANES) [15], applied to cuprates [16, 17] have shown a short-range atomic lattice reconstruction in cuprates below T^* with the appearance of an incommensurate modulation of local lattice distortions. This incommensurate lattice modulation of the CuO_2 plane [19] is related with the lattice misfit strain between layers [20, 21]. The strain field in cuprates is measured by the contraction of the Cu-O bond distance from the equilibrium distance of 197 pm [22] detected by XANES and EXAFS. In fact, pseudo Jahn Teller polarons show a self organisation above a critical misfit strain [23, 24]. The 1D lattice modulation of the superconducting planes has been proposed to induce a reconstruction of the 2D Fermi surface due to a periodic potential of 1D potential barriers in the superconducting planes in cuprates [28, 29] and in pnictides [25]. Moreover, experimental evidences are accumulating for a 2.5 Lifshitz transition associated with a vanishing Fermi surface in the case of iron based superconductors [30–33], diborides [26] and electron doped cuprates [34].

Moreover the physics of cuprates is controlled by an essential structural inhomogeneity [35, 36] that recently has been proposed to be made of scale invariant networks of




FIG. 1. The doping dependent isotope coefficient α (filled squares) and the critical temperature (open circles) of $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ (M=Sr in the lower panel, M=Ba in the upper panel) as a function of hole doping δ [44, 45]. The isotope coefficient for T_c is measured by oxygen isotope substitution, by replacing ^{16}O with ^{18}O and shows a pronounced maximum for both compounds near doping 1/8.

superconducting grains [37, 38]. This scenario has been recently observed by imaging lattice fluctuations using nano x-ray diffraction [18, 39, 40], where each superconducting grain of the network is tuned near the shape resonance in superconducting gaps [41].

The small Fermi surface pockets in $\text{YBa}_2\text{CuO}_{6+y}$ [1, 3, 4] are characterised by a very low value of the Fermi energy, of the order of 35-40 meV. Remarkably, this very small Fermi energy E_F is of the order of the maximum value of the superconducting gap Δ measured at low temperature below T_c ($\Delta/E_F \approx 1$) by angle resolved photo-emission spectroscopy (ARPES) and scanning tunnelling microscopy (STM).

The isotope coefficient α of the superconducting critical temperature is close to 0.5 in conventional BCS superconductors [42] and independent on the shift of the chemical potential. This result depends on the assumption of a single Fermi surface where the Fermi level is far from the band edge and the pairing attractive mechanism is the electron-phonon Cooper pairing. The experiments clearly show that this is not the case in cuprate high-temperature superconductors, where the isotope coefficient is doping dependent and nearly zero at optimum doping [43–50, 52–54]. This almost vanishing value of α has been considered a key evidence for an unconventional non-phononic pairing mechanism in high-temperature superconductors [51, 55]. Here we focus our interest on the available experimental isotope effect in $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ (M=Sr, Ba) [43–50, 52–54] since they show a single superconducting layer with the stripe phase at 1/8. On the contrary Y123 is a more complex bilayer cuprate

with complex organization of broken oxygen chains in the basal plane [53]. The isotope coefficient behavior in $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ (M=Sr, Ba) has a very particular doping dependence in $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ (M=Sr, Ba) [43–47], as shown in Fig. 1. It exhibits a strong anomaly clearly shown near doping 1/8 in Fig. 1, where the isotope coefficient α peak reaches a value close or larger than 0.5. In these in $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ systems the stripe phase is well established to appear at 1/8 doping. The large isotope effect supports the involvement of the lattice dynamics in the pairing, but in a non trivial way. The problem is very complex since there are also effects of the isotope substitution on the electronic structure [56] and on the stripe phase [17]. The isotope coefficient α is expected to increase near the insulator-to-metal transition, where the polaron scenario is dominant at the insulator-to-metal transition [77], but the sharp anomaly at a particular doping indicates that the isotope coefficient shows a different anomaly related with the metallic stripe phase. It has also been shown that α increases. The isotope effect has been discussed near a van Hove singularity in a single 2D band [57, 58] and in the frame of multi-band or multi-gap superconductivity far from band edges [59, 60], but there are missing theoretical efforts on the investigation of the anomalous isotope coefficient in a multi-band superconductor near a 2.5 Lifshitz transition at a metal-to-metal transition.

Theory. Multi-band multi-condensate superconductivity [60–63] is usually considered for two coupled BCS condensates where in the normal phase the Fermi energy is far from all band edges. The shape resonance in the superconducting gaps [61] is a type of Fano resonance between different pairing channels that occurs in multi-band metals where the chemical potential is driven at a 2.5 Lifshitz transition. The 2.5 Lifshitz transition in the proximity of a vanishing Fermi surface in a multi-band metal has been widely studied in the physics of Fermi surface topology in metals [64–67]. The shape resonances in superconducting gaps have been shown to occur a single 2D ultra-thin metallic layer [68] and in a metallic stripe with 1D sub-bands or mini-bands [69, 70]. In 3D multilayer materials [26, 27, 71, 72], and in a superlattice of stripes [28, 29] the superlattice reduces quantum fluctuations in low dimensions and the high T_c coherent phase can be realised. The T_c amplification is controlled by the Lifshitz energy parameter, measuring the energy difference between the chemical potential and the 2.5 Lifshitz transition. The maximum T_c is reached where the Lifshitz energy parameter is of the order of the energy cut-off for the pairing interaction [26, 27, 61].

In this work, we propose a model of a superlattice of quantum stripes that grabs the physical scenario emerging from the recent experiments discussed above. In fact, this model provides the appearing of a small electron pocket at the 2.5 Lifshitz transition as in cuprates. The fraction of the superconducting condensate originated by

this small pocket has a quasi bosonic character and it is located in the crossover regime of the BCS-BEC (Bose-Einstein Condensation) crossover, a phenomenon which is deeply studied in ultracold fermions [73–75].

Therefore, our model can reproduce the formation of a quasi-bosonic condensate in the phase space where the small pocket appears. This type of BCS-BEC crossover is a generic feature of multi-band / multi-condensate superconductors when the chemical potential is tuned close to the bottom of one of the bands and the pairing is strong enough to open gaps of the order of the (small) Fermi energy.

While the standard BCS-BEC crossover [76, 77] has been studied in single band metals, the present model provides a new scenario where the BCS-BEC crossover occurs in a multi-gap superconductor. Here, a first BCS condensate with order parameter Δ_1 in a large Fermi surface coexists with a second Bose-like condensate with order parameter Δ_2 in a second small electron pocket. Pair fluctuations and their screening in the multi-gap and multi-band (or multi-patch) models have been discussed also in the context of the physics of cuprates [78, 79].

In our model the Josephson-like coupling between the two condensates is a contact non retarded interaction due to pair exchange mechanisms, that can be attractive or repulsive. The intraband pairing is mediated by an effective attraction, having the momentum and energy structure of the electron-phonon interaction in the BCS approximation. Note that in the case of electronic mechanisms, such as exchange of spin fluctuations (paramagnons), the repulsive interaction transforms in an attractive pairing thanks to the d-wave symmetry of the superconducting order parameter, because the characteristic (large) wave-vector of the paramagnon connects state on the Fermi surface with opposite sign of the order parameter. Therefore both phononic and electronic mechanisms can be included in the model by an effective attractive interaction. This work is aimed to show that our simple theoretical model can explain the *doping dependent isotope coefficient* in cuprates near 1/8 doping in the framework of the striped phase, where the Fermi level is tuned near the 2.5 Lifshitz transition.

MODEL AND METHODS

Our model is based on two well established experimental facts for the physics near 1/8 doping in cuprates : i) a striped phase inducing Fermi surface reconstruction and 2) a 2.5 Lifshitz transition [1, 3, 4]. In the underdoped regime the cuprate superconductors show at doping 1/16 a first insulator-to-metal phase transition from the doped insulating Mott phase to a correlated striped metal and at doping 1/8 a metal-to-metal Lifshitz transition for the appearance or disappearance of a small electron Fermi surface pocket coexisting with Fermi arcs. To

investigate the response of the isotope effect at the Lifshitz transition, we consider the simplest physical model that grabs the essential physics of the cuprate metallic phase in the doping range near 1/8 doping: a superlattice of metallic stripes separated by a potential barrier [16, 19, 23, 24, 28, 69, 71] that makes a periodic potential in the 2D metallic layer, which provides the source for the Fermi surface reconstruction:

$$W(y) = \sum_{n=-\infty}^{+\infty} W_b(y - nl_p), \quad (1)$$

where $W_b(y) = -V_b$ for $|y| \leq L/2$ and $W_b(y) = 0$ for $L/2 < |y| < l_p/2$ where L is the width of the confining well and l_p is the periodicity of the superlattice in the y direction.

The confining potential of Eq.(1) generates a band structure organised in mini-bands. This model allows us to simulate the electronic structure of cuprates near the 2.5 Lifshitz transition for the appearing of a 2D FS. The Fermi surface in cuprates is simulated by tuning the Fermi level E_F below the bottom of the second superlattice mini-band E_2 where the superlattice FS (see Fig. 2) is made of a single FS of 1D character: the two open corrugated lines. The Fermi surface in cuprates is simulated by tuning the Fermi level E_F above the bottom of the second superlattice mini-band E_2 , where a second closed FS of 2D character appears beyond the first 1D mini-band (see Fig. 2). This is determined by the quasi free electron model dispersion of the second mini-band $E_{2,k}^D = E_2 + E(k_y) + k_x^2/2m$, where $E(k_y)$ is the energy dispersion in the y direction of the periodic potential $W(y)$ of the superlattice (here and in the following the reduced Planck constant is set to unity). The second closed Fermi surface changes its 2D topology into a 1D topology with isoenergetic open corrugated lines in the x direction, above some energy threshold $E_{2D-1D} - E_2 = \xi$, where ξ is the energy band dispersion in the y direction and E_{2D-1D} is the energy where the topology of the FS changes from 2D to 1D. Therefore in our model we tune the Fermi level in the energy range $E_2 < E_F < E_{2D-1D}$. Multi-Gap superconductivity in the energy range $E_2 - \xi < E_F < E_2 + \xi$ requires the theoretical approach we have recently proposed for diborides [26] and Fe-based [27] superconductors, which is capable to go beyond the standard BCS approximations, which consist of a single band, a large Fermi surface, a high Fermi energy, and a constant density of states (DOS) above and below E_F .

We consider a 1D periodic potential barrier of width B and wells of width L in the y direction with periodicity $l_p = L + B = 1.9nm$ to simulate the superstructure of Y-based and La-based cuprates and constant along the x direction. The potential barrier is fixed at $V_b = 1400meV$. We note that small variations of l_p and of other parameters of the potential do not influence the main results of




FIG. 2. The evolution of the Fermi surface (FS) in our model for the striped phase that simulates a 2.5 Lifshitz phase transition (P_L) at doping $\delta = P_L = 1/8$ in cuprates where a new detached electron-like Fermi surface appears. The inset of the left panel FS shows the lattice structure of a superlattice of stripes with finite hopping probability between stripes. E_c is the energy of the bottom of the second mini-band. The chemical potential is tuned from below (left panel $\delta < P_L$) to above (central panel $\delta > P_L$) where a new small 2D closed electron Fermi surface appears (red circle). The right panel shows the FS beyond the electronic topological transition (ETT) of the second band from a 2D FS to a corrugated 1D FS at higher doping called 2D-1D ETT.

this work. The strength of the potential barrier V_b is important to determine the 1D to 2D dimensional crossover observed in the phase diagram of cuprates: lower values of V_b will give a quasi-2D electronic system with no interesting stripe features, while larger values of V_b will lead to a strong 1D stripe physics, without the possibility of formation of electron pockets in the Brillouin zone, and without the screening of order parameter fluctuations active in a dimensional crossover (pronounced 1D anisotropy suppresses in general superconducting long-range order). The present choice of the periodic potential gives a band dispersion $\xi = 50\text{meV}$. The band dispersion ξ is two times the electronic hopping integral t_y in the direction y , that is much smaller than the hopping integrals t_x in the x direction. Solving the Schrödinger equation for the 1D periodic potential barrier of Eq.(1), we obtain the wave-functions of the electrons with a free electron band dispersion along the stripe direction and tight-binding mini-bands in the transverse direction. The eigenvalues are labelled by three quantum number $E = \epsilon_{n,k_x,k_y}$ where n is the mini-band index, k_x and k_y are the components of the electron wave-vectors in the superlattice. The DOS as function of the energy shows a jump at $E_{edge} = E_2$ and a sharp peak at E_{2D-1D} . The superconducting phase occurs because of the presence of an attractive intraband electron-electron effective interactions (1,1) and (2,2) in the first and second band respectively and the interband exchange-like interactions (1,2) and (2,1). The cutoff energies of the interactions are symmetrically fixed around the Fermi surface and the value of the effective coupling has been fixed at $\lambda = 1/3$. In the BCS approximation, i.e., a separable interaction in wave-vector space, the gap parameter has the same energy cut off as the interaction. Therefore it has a value

Δ_{n,k_y} around the Fermi surface in a range of energies equals to the energy cutoff, depending from the mini-band index n and the superlattice wave-vector k_y . The self consistent equation for the ground state ($T = 0$) energy gap Δ_{n,k_y} is:

$$\Delta_{n,k_y} = -\frac{1}{2N} \sum_{n',k'_y,k'_x} \frac{V_{\mathbf{k},\mathbf{k}'}^{n,n'} \Delta_{n',k'_y}}{\sqrt{(E_{n',k'_y} + \epsilon_{k'_x} - \mu)^2 + \Delta_{n',k'_y}^2}}, \quad (2)$$

where N is the total number of wave-vectors in the discrete summation, μ is the chemical potential, $V_{\mathbf{k},\mathbf{k}'}^{n,n'}$ is the effective pairing interaction

$$V_{\mathbf{k},\mathbf{k}'}^{n,n'} = \tilde{V}_{\mathbf{k},\mathbf{k}'}^{n,n'} \times \theta(\omega_0 - |E_{n,k_y} + \epsilon_{k_x} - \mu|) \theta(\omega_0 - |E_{n',k'_y} + \epsilon_{k'_x} - \mu|) \quad (3)$$

calculated taking into account the interference effects between the wave functions of the pairing electrons in the different mini-bands, where n and n' are the mini-band indexes, $k_y(k'_y)$ is the superlattice wave-vector and $k_x(k'_x)$ is the component of the wave-vector in the stripe direction of the initial (final) state in the pairing process, and

$$\tilde{V}_{\mathbf{k},\mathbf{k}'}^{n,n'} = -\frac{\lambda_{n,n'}}{N_0} S \times \int_S \psi_{n',-k'_y}(y) \psi_{n,-k_y}(y) \psi_{n,k_y}(y) \psi_{n',k'_y}(y) dx dy, \quad (4)$$

where N_0 is the DOS at E_F for a free-electron 2D system, $\lambda_{n,n'}$ is the dimensionless coupling parameter, $S = L_x L_y$ is the surface of the plane and $\psi_{n,k_y}(y)$ are the eigenfunctions in the superlattice of quantum stripes. The gap equations have been solved iteratively. We obtain anisotropic gaps strongly dependent on the mini-band index and weakly dependent on the superlattice wave-vector k_y . According with Leggett [76], the ground-state BCS wave function corresponds to an ensemble of overlapping Cooper pairs at weak coupling (BCS regime) and evolves to molecular (non-overlapping) pairs with bosonic character. The point is that the BCS equation for the gap has to be coupled to the equation that fixes the fermion density: with increasing coupling (or decreasing density), the chemical potential μ results strongly normalised with respect to the Fermi energy E_F of the non interacting system, and approaches minus half of the molecular binding energy of the corresponding two-body problem in the vacuum. Therefore in order to correctly describe the case of the chemical potential near a band edge, where all electrons in the new appearing band condense forming a bosonic-like gas in the second mini-band,

the chemical potential in the superconducting phase is normalised by the gaps opening at any chosen value of the charge density ρ :

$$\begin{aligned} \rho &= \frac{1}{L_x L_y} \sum_n \sum_{k_x, k_y} \left[1 - \frac{E_{n, k_y} + \epsilon_{k_x} - \mu}{\sqrt{(E_{n, k_y} + \epsilon_{k_x} - \mu)^2 + \Delta_{n, k_y}^2}} \right] \\ &= \frac{\delta k_y}{\pi} \sum_{n=1}^{N_b} \sum_{k_y=0}^{\pi/l_p} \int_0^{\epsilon_{min}} d\epsilon \frac{2N(\epsilon)}{L_x} + \int_{\epsilon_{min}}^{\epsilon_{max}} d\epsilon \frac{N(\epsilon)}{L_x} \\ &\times \left(1 - \frac{E_{n, k_y} + \epsilon_{k_x} - \mu}{\sqrt{(E_{n, k_y} + \epsilon_{k_x} - \mu)^2 + \Delta_{n, k_y}^2}} \right), \end{aligned} \quad (5)$$

where

$$\begin{aligned} \epsilon_{min} &= \max[0, \mu - \omega_0 - E_{n, k_y}], \\ \epsilon_{max} &= \max[0, \mu + \omega_0 - E_{n, k_y}], \\ N(\epsilon) &= \frac{L_x}{2\pi\sqrt{\frac{\epsilon}{2m}}}, \end{aligned}$$

and N_b is the number of the occupied mini-bands, L_x and L_y are the size of the considered surface and the increment in k_y is taken as $\delta k_y = 2\pi/L_y$. We compute the critical temperature T_c of the superconducting transition solving the linearised BCS equations

$$\Delta_{n, k_y} = -\frac{1}{2N} \sum_{n', k'} V_{\mathbf{k}, \mathbf{k}'}^{n, n'} \frac{\tanh(\frac{E_{n, k_y} + \epsilon_{k_x} - \mu}{2T_c})}{E_{n, k_y} + \epsilon_{k_x} - \mu} \Delta_{n', k'}, \quad (6)$$

where the energy dispersion is measured with respect to the chemical potential. The iterations are stopped when a convergence factor of 10^{-6} has been reached, starting with a trial temperature T_1 and finding the T_c by the Newton tangent method to solve the implicit integral equation for T_c . The T_c is evaluated as a function of the chemical potential in the proximity of the edge of the 2-nd mini-band. The tuning of the chemical potential is measured by the Lifshitz parameter $z = (\mu - E_2)/\omega_0$ where E_2 is the bottom of the second band and ω_0 is the energy cut-off for the pairing interaction.

THE ISOTOPE COEFFICIENT

The experimental 2.5 Lifshitz phase transition observed in the experiment reported by ref. [1–3] at doping $\delta = P_L = 1/8$ (reduced doping $\nu = 16\delta - 2 = 0$) is simulated by tuning the Fermi level at $E_F = E_2$, *i.e.*, at $z = 0$, corresponding with the reduced doping range $-1 < \nu < +1$. The gaps, the T_c and the gap to T_c ratios have been plotted as a function of the Lifshitz energy parameter in the range $-1 < z < +1$, as shown in Figure 3.




FIG. 3. a) The superconducting gaps in the first and second mini-bands normalised to the maximum value of the gap in the second mini-band; b) the critical temperature T_c normalised to its maximum value; c) and the gaps to T_c ratios for the two gaps as a function of the Lifshitz parameter $z = (\mu - E_2)/\omega_0$ compared with the standard BCS value of 3.5.

We obtain the minimum of T_c where μ is tuned at $z = 0$, *i.e.*, at the 2.5 Lifshitz transition for the appearance of a new detached electron Fermi surface that well reproduce the minimum at $1/8$ in Fig. 1. The maximum of T_c occurs where the topology of the FS of the 2-nd sub-band shows the 2D- $\rightarrow$ 1D electronic topological transition. This result is assigned to the shape resonance in the two superconducting gaps controlled by the exchange-like pair-transfer (or Josephson-like) pairing that shows a minimum (maximum) of T_c at the antiresonance at $z = -1$ (resonance at $z = 1$) due to the negative (positive) quantum interference effects typical of the shape resonances in the superconducting gaps.

Our results well reproduce the increase of the critical temperature T_c going from 0.125 to 0.18 doping in cuprates. The isotope effect $\alpha = \partial \ln T_c / \partial \ln M$ is calculated with the assumption of an energy cutoff of the interaction dependent from the isotopic mass as $\omega \propto M^{-1/2}$.

In Fig. 4 we report the calculated isotopic coefficient α as a function of the chemical potential that strongly deviates from the standard BCS value $\alpha = 0.5$. We find the maximum of T_c and the minimum of α at the shape resonance $\mu = E_{2D-1D}$, because the T_c is weakly dependent by small variation of the energy cutoff. On the contrary by tuning the chemical potential at the 2.5 Lifshitz transition for the appearance of the second circular Fermi surface, *i.e.*, at the band edge of the second miniband $\mu = E_2$, we find a large value of $\alpha \gg 0.5$ and a drop of T_c in agreement with the experimental data. The isotope coefficient has been calculated considering the cases where only one of the intraband pairing energies $\omega_{11} \propto M^{-1/2}$ or $\omega_{22} \propto M^{-1/2}$ and the interband pairing energy is isotope dependent or independent. The best agreement with the experimental data reported in Fig. 4 is obtained for the case where both intraband pairing energy $\omega_{11} \propto M^{1/2}$ and $\omega_{22} \propto M^{1/2}$ are isotope dependent and the electronic interband energy is isotope independent. Note that at the Lifshitz phase transition at ($z = 0$) both α and the gap to T_c ratios in both bands get exactly the conventional BCS values. This is the crossing point between the Bose-like regime where Δ_2 is smaller than Δ_1 but larger than $\mu - E_2$ (*i.e.*, the Fermi surface in band 2 is destroyed by the gap opening) and the BCS-like regime where Δ_2 is larger than Δ_1 but smaller than $\mu - E_2$ (*i.e.*, the Fermi surface in band 2 is only partially smeared by the gap opening).

DISCUSSIONS AND CONCLUSIONS

Starting from the stripes scenario near 1/8 for the electronic structure of cuprate superconductors and the identification of the 2.5 Lifshitz transition in the phase diagram of cuprates, we study the behaviour of the critical temperature T_c and the isotope coefficient of the superconducting phase transition as a function of the chemical potential near the 2.5 Lifshitz metal-to-metal transition, which is located at the band edge of a sub-band for a superlattice of quantum stripes.

We obtain an interesting asymmetric feature of the chemical potential dependence of T_c and α . This peculiar shapes of T_c and α given by our numerical calculations are well in agreement with the phenomenology of cuprate superconductors near doping 1/8. The energy cutoff ω_0 of the effective pairing interaction considered in our model determines the width of the shape resonance in the superconducting gaps and the small value of the isotope coefficient in the flat region of its doping dependence, well below the standard BCS value of $\alpha=0.5$, where T_c has a maximum in agreement with experiments. We find the maximum T_c when the chemical potential is tuned near the 2D-1D electronic topological transition of the 2-nd sub-band (where the Lifshitz parameter assumes the value $z = 1$). The maximum value




FIG. 4. panel a: The experimental isotope coefficient α as a function of the reduced doping $\nu = 16\delta - 2$ for $La_{2-x}M_xCuO_4$ ($M = Sr$ dots and $M = Ba$ squares) compared with the theoretical isotope coefficient α as a function of the Lifshitz parameter for the case of an isotope effect (IE) driven by intraband pairing in both bands $\omega_{11} \propto M^{1/2}$ and $\omega_{22} \propto M^{1/2}$ while the Josephson-like exchange-like pair transfer mechanism is isotope independent; panel b) theoretical isotope coefficient α as a function of the Lifshitz parameter for the case of an isotope effect (IE) only the intraband pairing energy in band 1 $\omega_{11} \propto M^{1/2}$ (filled dots) and for the case of an isotope effect (IE) only the intraband pairing energy in band 2 $\omega_{22} \propto M^{1/2}$ (empty triangles).

of the isotope coefficient is reached at the 2.5 Lifshitz transition, in the range of the Fano antiresonance where T_c is strongly suppressed. Interestingly, the ratios between the gaps and T_c in different sub-bands cross the conventional BCS ratio values ($=3.5$) at the 2.5 Lifshitz transition, while sizable deviations from the conventional value are obtained in our calculations above this transition, in the range $1 < z < 2$ accessible to experiments.

Therefore, we have provided a simple model that reproduces the enhancing of the isotope coefficient well above the conventional BCS value and explains the rapid doping dependence of the isotope coefficient α observed in the $La_{2-x}M_xCuO_4$ (La214) and $YBa_2Cu_3O_{6+y}$ (Y123) systems near doping 1/8. The present results have an impact on the physics of superconductivity in nano-sized superconductors [51, 55, 70], where the shape resonance in superconducting gaps is gaining momentum as a key ingredient for the road map for new high- T_c superconductors. Moreover, shape resonances and quantum size

effects have been also considered in the context of superconductivity in nanofilms [80] and superfluidity in cigar-shaped ultracold Fermi gases [81] as a possible new driving mechanism to tune an atypical BCS-BEC crossover in multi-band fermionic systems, with the appearance of a coherent mixture of BCS-like and BEC-like condensate. Finally, we have shown that the anomalous maximum of the isotope effect and its peculiar doping dependence in the doping range $1/8$, shown in Fig. 1 in La₂14 families, could have its origin in a 2.5 Lifshitz transition for a metal-to-metal transition in a multi-band multi-condensate superconductor, for the appearing or disappearing of small electron-like Fermi surface pocket. Therefore, the experimental investigation of the complex doping dependent isotope coefficient α in the $1/8$ doping range in other cuprate and pnictides families of superconductors could provide a signature for the role of the 2.5 Lifshitz transitions in the superconducting phase, in agreement with a large Nernst effect measured in the normal phase [65, 66].

-
- [1] D. LeBoeuf, N. D. Leyraud, B. Vignolle, M. Sutherland, B. J. Ramshaw, J. Levallois, R. Daou, F. Laliberte, O. C. Choinire, J. Chang *et al.*, Phys. Rev. B **83**, 054506 (2011).
 - [2] S. E. Sebastian, N. Harrison and G. G. Lonzarich Phil. Trans. R. Soc. A **369**, 1687-1711 (2011).
 - [3] F. Laliberte, J. Chang, N. Doiron-Leyraud, E. Hassinger, R. Daou, M. Rondeau, B. J. Ramshaw, R. Liang, D. A. Bonn, W. N. Hardy, S. Pyon, T. Takayama, H. Takagi, I. Sheikin, L. Malone, C. Proust, K. Behnia, and L. Taillefer, Nature Communications **2**, 432 (2011).
 - [4] R. Khasanov, S. Strassler, D. Di Castro, T. Masui, S. Miyasaka, S. Tajima, A. Bussman-Holder, and H. Keller, Phys. Rev. Lett. **99**, 237601 (2007).
 - [5] M. C. Boyer, W. D. Wise, K. Chatterjee, M. Yi, T. Kondo, T. Takeuchi, H. Ikuta, and E. W. Hudson, Nature Physics **3**, 802 (2007).
 - [6] S. S. Botelho and C. A. R. Sá de Melo, Phys. Rev. B **71**, 134507 (2005).
 - [7] S. G. Ovchinnikov, M. M. Korshunov, and E. I. Shneyder, Journal of Experimental and Theoretical Physics **109**, 5775 (2009).
 - [8] S. Sakai, Y. Motome, and M. Imada, Phys. Rev. Lett. **102**, 056404 (2009).
 - [9] M. R. Norman and J. Lin, and A. J. Millis Phys. Rev. B **81**, 180513 (2010).
 - [10] M. Norman, Physics **3**, 86 (2010).
 - [11] A. Chubukov, Physics **3**, 54 (2010).
 - [12] R. J. Birgeneau, C. Stock, J. M. Tranquada, and K. Yamada, Journal of the Physical Society of Japan **75**, 111003 (2006).
 - [13] S. B. Wilkins, M. P. M. Dean, J. Fink, M. Huecker, J. Geck, V. Soltwisch, E. Schierle, E. Weschke, G. Gu, S. Uchida, N. Ichikawa, J. M. Tranquada, and J. P. Hill, Phys. Rev. B **84**, 195101 (2011).
 - [14] A. Bianconi, R. Z. Bachrach, Phys. Rev. Lett. **42**, 104 (1979).
 - [15] A. Bianconi, R. S. Bauer, Surface Science **99**, 76-86 (1980).
 - [16] A. Bianconi, N.L. Saini, A. Lanzara, M. Missori, T. Rossetti, H. Oyanagi, H. Yamaguchi, K. Oka and T. Ito, Phys. Rev. Lett. **76**, 3412 (1996).
 - [17] A. Lanzara, Guo-meng Zhao, N. L. Saini, A. Bianconi, K. Conder, H. Keller, and K. A. Müller J. Phys. :Condens. Matter **11**, 541 (1999).
 - [18] N. Poccia, A. Ricci, G. Campi, M. Fratini, A. Puri, D. Di Gioacchino, A. Marcelli, M. Reynolds, M. Burghammer, N. L. Saini, G. Aeppli, A. Bianconi Proc. Natl. Acad. Sci. USA **109** (2012) doi:10.1073/pnas.1208492109 ; preprint arXiv:1208.0101
 - [19] A. Bianconi, N. L. Saini, T. Rossetti, A. Lanzara, A. Perali, M. Missori, H. Oyanagi, H. Yamaguchi, Y. Nishihara, and D. H. Ha, Phys. Rev. B **54**, 12018 (1996).
 - [20] A. Bianconi, S. Agrestini, G. Bianconi, D. Di Castro, N. L. Saini, Journal of alloys and compounds **317**, 537 (2001).
 - [21] S. Agrestini, N. L. Saini, G. Bianconi, A. Bianconi, Journal of Physics A: Mathematical and General **36**, 9133 (2003).
 - [22] J. Garcia, M. Benfatto, C. R. Natoli, A. Bianconi, A. Fontaine, H. Tolentino Chemical Physics **132**, 295 (1989).
 - [23] A. Bianconi, Solid State Communications **91**, 1 (1994).
 - [24] A. Bianconi, M. Missori, Solid State Communications **91**, 287 (1994).
 - [25] R. Caivano, M. Fratini, N. Poccia, A. Ricci, A. Puri, Z.-A. Ren, X.-L. Dong, J. Yang, W. Lu, Z.-X. Zhao, L. Barba and A. Bianconi Supercond. Sci. Technol. **22**, 014004 (2009).
 - [26] D. Innocenti, N. Poccia, A. Ricci, A. Valletta, S. Caprara, A. Perali, and A. Bianconi, Phys. Rev. B **82**, 184528 (2010).
 - [27] D. Innocenti, S. Caprara, N. Poccia, A. Ricci, A. Valletta, and A. Bianconi, Supercond. Sci. Technol. **24**, 015012 (2011).
 - [28] A. Bianconi, A. Valletta, A. Perali, and N. L. Saini, Solid State Communications **102**, 369 (1997).
 - [29] A. Bianconi, A. Valletta, A. Perali, N. L. Saini, Physica C **296**, 269 (1998).
 - [30] Chang Liu, Takeshi Kondo, Rafael M. Fernandes, Ari D. Palczewski, Eun Deok Mun, Ni Ni, Alexander N. Thaler, Aaron Bostwick, Eli Rotenberg, Jörg Schmalian, Sergey L. Bud'ko, Paul C. Canfield and Adam Kaminski, Nature Physics **6**, 419 (2010).
 - [31] C. Liu, A. D. Palczewski, R. S. Dhaka, T. Kondo, R. M. Fernandes, E. D. Mun, H. Hodovanets, A. N. Thaler, J. Schmalian, S. L. Bud'ko, P. C. Canfield, and A. Kaminski, Phys. Rev. B **84**, 020509 (2011).
 - [32] A. A. Kordyuk, V. B. Zabolotnyy, D. V. Evtushinsky, A. N. Yaresko, B. Büchner, S. V. Borisenko, preprint arXiv:1111.0288 (2011).
 - [33] S.V. Borisenko, V. B. Zabolotnyy, A. A. Kordyuk, D. V. Evtushinsky, T. K. Kim, I. V. Morozov, R. Follath, B. Büchner, Symmetry **4**, 251-264 2012; doi:10.3390/sym4010251 (arXiv:1110.6922 (2011)).
 - [34] M. V. Kartsovnik, T. Helm, C. Putzke, F. Wolff-Fabris, I. Sheikin, S. Lepault, C. Proust, D. Vignolles, N. Bittner, W. Biberacher, A. Erb, J. Wosnitzer, and R. Gross, New J. Phys. **13**, 015001 (2011).
 - [35] E. Dagotto, Science **309**, 257 (2005).
 - [36] K. A. Müller, Journal of Physics: Condensed Matter **19**,

- 251002 (2007).
- [37] G. Bianconi, Phys. Rev. E **85**, 061113 (2012).
 - [38] G. Bianconi, Journal of Statistical Mechanics: Theory and Experiment **2012**, P07021 (2012).
 - [39] M. Fratini, N. Poccia, A. Ricci, G. Campi, M. Burghammer, G. Aeppli, A. Bianconi Nature **466**, 841 (2010).
 - [40] N. Poccia, M. Fratini, A. Ricci, G. Campi, L. Barba, A. Vittorini-Orgeas, G. Bianconi, G. Aeppli, A. Bianconi, Nature Materials **10**, 733 (2011).
 - [41] A. Bianconi, D. Di Castro, N. L. Saini, and G. Bianconi, in Phase Transitions and Self-Organization in Electronic and Molecular Networks, edited by M. F. Thorpe and J. C. Phillips (Kluwer Boston, 2002), Fundamental Materials Research, chap. 24, pp. 375-388.
 - [42] E. Maxwell, Phys. Rev. **78**, 477 (1950).
 - [43] Y. Sassa *et al.*, Phys. Rev. B **83**, 140511 (2011).
 - [44] M. K. Crawford, M. N. Kunchur, W. E. Farneth, E. M. McCarron, and S. J. Poon, Phys. Rev. B **41**, 282 (1990).
 - [45] M. K. Crawford, W. E. Farneth, E. M. McCarron, R. L. Harlow, and A. H. Moudden, Science **250**, 1390 (1990).
 - [46] G.-M. Zhao, M. B. Hunt, H. Keller, and K. A. Müller, Nature **385**, 236 (1997).
 - [47] J. P. Franck, S. Harker, and J. H. Brewer, Phys. Rev. Lett. **71**, 283 (1993).
 - [48] J. P. Franck, J. Jung, Mohamed, S. Gyax, and G. I. Sproule, Phys. Rev. B **44**, 5318 (1991).
 - [49] R. Khasanov, A. Shengelaya, D. Di Castro, E. Morenzoni, A. Maisuradze, I. M. Savi, K. Conder, E. Pomjakushina, A. Bussmann-Holder, and H. Keller, Phys. Rev. Lett. **101**, 077001 (2008).
 - [50] J. P. Franck, in Physical Properties of High Temp Superconductors IV, edited by D.M. Ginzberg (World Sci., Singapore, 1994) pp 189-293.
 - [51] D. Zech, H. Keller, K. Conder, E. Kaldis, E. Liarokapis, N. Poulakis, and K. A. Müller, Nature **371**, 681 (1994).
 - [52] D. Zech, K. Conder, H. Keller, E. Kaldis, and K. A. Müller, Physica B **219-220**, 136-138 (1996).
 - [53] H. Keller and A. Bussmann-Holder, Advances in Condensed Matter Physics **2010**, 393526 (2010).
 - [54] X. J. Chen, V. V. Struzhkin, Z. Wu, H. Q. Lin, R. J. Hemley, and H. K. Mao, Proceedings of the National Academy of Sciences of the United States of America **104**, 3732 (2007).
 - [55] V. Z. Kresin, A. Bill, S. A. Wolf, and Yu, Phys. Rev. B **56**, 107 (1997).
 - [56] A. Lanzara *et al.* Nature **412**, 510-514 (2001).
 - [57] R. Markiewicz, Physica C: Superconductivity **153-155**, 1181 (1988).
 - [58] C. C. Tsuei, D. M. Newns, C. C. Chi, and P. C. Pattnaik, Phys. Rev. Lett. **65**, 2724 (1990).
 - [59] P. Konsin, N. Kristoffel, and T. Örd, Annalen der Physik **505**, 279 (1993).
 - [60] A. Bussmann-Holder, H. Keller, R. Khasanov, A. Simon, A. Bianconi, and A. R. Bishop, New Journal of Physics **13**, 093009 (2011).
 - [61] A. Bianconi, Journal of Superconductivity **18**, 625 (2005).
 - [62] N. Kristoffel, P. Rubin, and T. Ord, Journal of Superconductivity and Novel Magnetism **20**, 571 (2007).
 - [63] A. A. Golubov and A. E. Koshelev Phys. Rev. B **83**, 094521 (2011).
 - [64] I. M. Lifshitz, Zh. Eksp. Teor. Fiz. **38**, 1569 (1960); and in English in [Sov. Phys. JETP **11**, 1130 (1960)].
 - [65] Ya. M. Blanter, M. I. Kaganov, A.V. Pantsulaya, A. A. Varlamov, Physics Reports **245**, 159 (1994).
 - [66] A. E. Koshelev, A. A. Varlamov, and V. M. Vinokur Phys. Rev. B **72**, 064523 (2005).
 - [67] A. Levchenko, M. R. Norman, and A. A. Varlamov Phys. Rev. B **83**, 020506 (2011).
 - [68] J. M. Blatt and C. J. Thompson, Phys. Rev. Lett. **10**, 332 (1963).
 - [69] A. Perali, A. Bianconi, A. Lanzara, N. L. Saini, Solid State Communications **100**, 181 (1996).
 - [70] A. A. Shanenko and M. D. Croitoru, Phys. Rev. B **73**, 012510 (2006).
 - [71] A. Bianconi, Solid State Communications **89**, 933 (1994).
 - [72] A. Bussmann-Holder and A. Bianconi Phys. Rev. B **67**, 132509 (2003).
 - [73] J. P. Gaebler *et al.*, Nature Phys. **6**, 569 (2010).
 - [74] A. Perali *et al.*, Phys. Rev. Lett. **106**, 060402 (2011).
 - [75] F. Palestini, A. Perali, P. Pieri, and G. C. Strinati, Phys. Rev. B **85**, 024517 (2012).
 - [76] A. J. Leggett, in Modern Trends in the Theory of Condensed Matter, edited by A. Pekalski and R. Przys-tawa, Lecture Notes in Physics Vol. **115** (Springer-Verlag, Berlin, 1980), p. 13.
 - [77] A. S. Alexandrov, Phys. Rev. B **46**, 14932 (1992).
 - [78] A. Perali, C. Castellani, C. Di Castro, M. Grilli, E. Piegari, and A. A. Varlamov, Phys. Rev. B **62**, 9295 (2000).
 - [79] A. Perali, M. Sindel, and G. Kotliar, Eur. Phys. J. B **24**, 487 (2001). A. Malinowski *et al.*, Phys. Rev. B **66**, 104512 (2002).
 - [80] Y. Chen, A. A. Shanenko, A. Perali and F. M. Peeters, J. Phys.: Condens. Matter **24**, 185701 (2012).
 - [81] A. A. Shanenko, M. D. Croitoru, A. V. Vagov, V. M. Axt, A. Perali, and F. M. Peeters, Phys. Rev. A (2012). arXiv:1203.3325